

LOW MASS EXOTIC BARYONS: MYTH OR REALITY ?

B. Tatischeff

Institut de Physique Nucléaire, CNRS/IN2P3, F-91406 Orsay Cedex, France

E-mail: tati@ipno.in2p3.fr

Abstract

Using mainly $\bar{p} p \rightarrow p \pi^+ X$ and $\bar{p} p \rightarrow p_f p_s X$ reactions, narrow baryonic structures were observed in the mass range $950 \leq M \leq 1800$ MeV.

Key-words: experiment, narrow, exotic, baryons, multiquark, states

1 Introduction

Narrow structures in hadrons have been observed since two decades. Despite this long time these results are still in debate. Two reasons explain, if not justify, such situation. First the observed structures are often small, and some observations were made using bubble chamber slides, therefore with small statistics. The second reason is that their genuine existence constitute a disturbing element inside the present theories.

Narrow structures in baryons were first observed in the missing mass M_X of the $p p \rightarrow p \pi^+ X$ reaction, in the mass range $1.0 \leq M \leq 1.1$ GeV [1]. The two lowest structure masses are small, preventing them from disintegration through pion emission. When all three structures inside this mass range at $M=1004$ MeV, 1044 MeV, and 1094 MeV are observed at nearly all angles and energies, narrow structures are also observed in other mass ranges, but they are more weakly excited, and therefore extracted with less confidence.

The experiment was performed at the SPES3 beam line using the Saturne polarized proton beam ($T_p=1.52$ GeV, 1.805 GeV, and 2.1 GeV). Narrow baryons were observed in the missing mass M_X and in the invariant masses $M_{p\pi^+}$ and $M_{X\pi^+}$ of the $p p \rightarrow p \pi^+ X$ reaction, and in the invariant masses M_{Xp_f} and M_{Xp_s} of the $p p \rightarrow p_f p_s X$ reaction. Here p_f (p_s) means fast (slow) proton. Some structures were also observed at nearby masses in the missing mass of the $d p \rightarrow p p X$ reaction. Fig. 1 shows the mass ranges studied in the Saturne (SPES3) experiment. The three vertical strips for each variable correspond to the three incident energies, increasing from left to right. The horizontal lines show the masses where a narrow structure was observed.

Figure 1: Experimental ranges for the study of narrow baryonic masses (in GeV).

Figure 2: Selection of two cross sections showing an oscillatory pattern in the missing mass.

2 Narrow baryons in the mass range $950 \leq M \leq 995$ MeV

When the missing mass spectra display smooth patterns up to $M=950$ MeV, they display oscillatory patterns for larger masses. An example of such oscillations is shown in Fig. 2.

A systematic control was undertaken to make sure that these structures were not produced by the lost event corrections or by the acceptance renormalizations. Fig. 3 shows that the analyzing powers display - like the cross sections - a non smooth behavior. Here only small angle data are shown, since the error bars increase quickly for angles $\geq 9^\circ$.

Figure 3: Selection of small angle analyzing powers of the missing mass M_X from the $\vec{p} p \rightarrow p \pi^+ X$ reaction. The two columns correspond from left to right to $T_p=1520$ MeV and $T_p=1805$ MeV.

For the observed oscillatory shapes in cross sections, the background determination is clearly somewhat ambiguous. Several different choices can be used for background. The first one consists to draw an averaged background inside the data. Then all structures will be strongly reduced, if not disappear. However we consider as quit unlikely the situation where nearly all spectra will exhibit an oscillatory pattern and where all these oscillations will be accidental. The second choice consists to consider these variations as physical, and to give them the same width as the experimental neutron missing mass widths which increased slowly with the spectrometer angle. Then the structures will be extracted with a

higher number of standard deviation (S.D.). We choose the third intermediate choice and draw the background using the low data points as in Fig. 2.

Although 54 peaks were extracted from all cross section spectra, only 18 were kept which had a valuable statistical significance, since they were extracted with $S.D. \geq 3$. The corresponding masses are displayed in Fig. 4. When a mass (± 1 MeV) is obtained at least twice, it is considered as being a candidate for a new structure and an horizontal dashed range is drawn in Fig. 4. The figure shows also the two narrow masses (at 966 MeV and 986 MeV), extracted by L.V. Filkov *et al.* from the $p d \rightarrow p p X_1$ reaction [2].

Figure 4: Low narrow exotic baryonic masses.

We tentatively associate these structures to exotic baryons formed by two coloured quark clusters, and we use phenomenologically a mass formula derived 23 years ago [3]:

$$M = M_0 + M_1[i_1(i_1 + 1) + i_2(i_2 + 1) + (1/3)s_1(s_1 + 1) + (1/3)s_2(s_2 + 1)] \quad (1)$$

where $s_1(s_2)$ and $i_1(i_2)$ are the first (second) cluster spin and isospin values. The two parameters, $M_0=924$ MeV, and $M_1=15.6$ MeV are adjusted in order to describe the neutron mass and the first exotic mass at 950 MeV. The other masses, possible spins and isospins obtained with the use of this formula, are shown in Fig. 5. We observe the good correspondence between the experimental and the calculated masses.

Figure 5: Experimental and calculated exotic baryonic masses in the range $948 \leq M \leq 995$ MeV.

3 Narrow baryons in the mass range $1100 \leq M \leq 1400$ MeV

Some results corresponding to this mass range were already reported [4]. Narrow peaks were extracted from missing mass or invariant mass spectra. Only those structures observed several times at the same mass ± 5 MeV were taken into account. A selection of some spectra showing 4 peaks close to 1136 MeV is presented in Fig. 6.

Several structures were observed in this mass range, extracted from the experiments

Figure 6: Selection of four spectra illustrating a narrow structure close to $M_X=1136$ MeV.

quoted previously, and also from the invariant mass $M_{p\pi}$ of the $\gamma n \rightarrow \pi^0 \pi^- p$ reaction studied at MAMI [5], and from the center of mass energy of the $\gamma p \rightarrow \pi^+ n$ reaction studied at Bonn [6]. The results are all listed in Fig. 7 which is explained in table 1.

The same mass formula as the one discussed before, was used in this mass range. The two parameters $M_0=838.2$ MeV and $M_1=100.3$ MeV were adjusted in order to describe the mass (and possible spin and isospin) of the neutron and of the Roper $N(1440)$ MeV resonance. Fig. 8 shows the good agreement between the experimental and the calculated masses. The last column in Fig. 8 shows the masses calculated by N. Konno, H. Noya and H. Nakamura [7] inside their diquark cluster model.

Figure 7: Narrow structure baryonic masses observed in cross sections from different reactions in the range $1100 \leq M \leq 1400$ MeV.

4 Narrow baryons in the mass range $1500 \leq M \leq 1700$ MeV

Starting from this mass range, we expect to observe a complex situation, due to a superposition of weakly excited structures as those discussed previously with interferences from many broad overlapping PDG resonances having the same quantum numbers. Many such resonances overlap, as N^* , $J^P=1/2^-, 1/2^+, 3/2^+$, or Δ , $J^P=3/2^+, 1/2^-, 3/2^-$, and $1/2^+$ in the mass range studied here. We observe indeed structures slipping smoothly in mass for increasing spectrometer angles, superimposed to structures stable in mass. The data analysis in this mass range is not completed.

5 Narrow baryons in the mass range $1720 \leq M \leq 1800$ MeV

Surprisingly, the superposition of smoothly moving structures with structures stable in mass (those we are looking for here) seems not impede us in this mass range. Three structures at $M=1727$ MeV, 1748 MeV, and 1771 MeV were observed at different angles, in different variables from both reactions using our highest incident proton energy: $T_p=2.1$ GeV. These masses were observed in ten different experimental conditions, always - but in one case - with $SD \geq 3$. Fig. 9 shows some illustrations of these results. The insert (d) shows an analyzing power data where narrow structures are present at the same masses where they appear in cross sections, namely at 1727 MeV and 1748 MeV.

6 Conclusion

Narrow, low mass, non strange structures were observed in baryonic missing masses and in baryonic invariant masses studied with help of different reactions. These weakly excited narrow structures were observed in the whole baryonic mass range investigated: $940 \leq M \leq 1800$ MeV.

Many checks were undertaken to make sure that these structures were not produced by experimental artifacts [8]. Their small widths and the stability of the extracted masses, regardless of the experiment, of the variable under study, of the incident energy, and of the reaction angle (see Fig. 1), allowed us to conclude that these structures are genuine and not produced by dynamical rescatterings. We concluded that these peaks correspond to new exotic baryons since there is no room for them within the many theoretical constituent quark models [9].

Figure 8: Calculated and experimental narrow exotic baryon level spectrum in the range $1100 \leq M \leq 1400$ MeV.

Figure 9: Selection of three cross sections and one analyzing power data showing narrow structures in the mass range $1720 \leq M \leq 1800$ MeV.

Several theoretical attempts were performed to explain these observations:

A.P. Kobushkin [10] assumes that the first narrow baryons ($M \leq 1075$ MeV) may be members of a total antisymmetric representation of the spin-flavor group.

N. Konno, H. Noya, and H. Nakamura [7] assume that the exotic baryons consist of a diquark and a quark (Diquark cluster model).


Y.E. Pokrovsky [11] derives a model which deals with chiral and scale symmetries of QCD, and describes hadrons as multi-bag states ($Bag\bar{Bag}Bag$) for baryons.



Th. Walcher [12] derives a model based on the excitation of quark condensate. The model explains the narrow states as a multiple production of a “genuine” Goldstone Boson with a mass close to 20 MeV.



We tentatively associate our narrow baryons with exotic baryons made of two coloured quark clusters [1] [4].


References





- [1] B. Tatischeff *et al.*, Phys. Rev. Lett. **79**, 601 (1997).
- [2] L.V. Fil’kov *et al.*, Eur. Phys. J. **A 12**, 369, (2001).
- [3] P.J. Mulders, A.T. Aerts, and J.J. de Swart, Phys. Rev. **D19**, 2635 (1979).
- [4] B. Tatischeff, Proceedings of the IX Int. Conf. on Hadron Spectroscopy, Protvino, august 2001, edited by D. Amelin, AIP Conf. Proc., **619**, p 765.
- [5] A. Zabrodin *et al.*, Phys. Rev. **C60**, 055201, (1999).
- [6] H.W. Dannhausen *et al.*, Eur. Phys. J. **A 11**, 441 (2001).
- [7] H. Noya and H. Nakamura, Proceedings of the IX Int. Conf. on Hadron Spectroscopy, Protvino, august 2001, edited by D. Amelin, AIP Conf. Proc., **619**, p 533.
- [8] B. Tatischeff *et al.*, to be published, and nucl-ex/0207003.
- [9] S. Capstick and W. Roberts, Prog. Part. Nucl. Phys. **45**, 0S241 (2000).
- [10] A.P. Kobushkin, nucl-th/9804069.
- [11] Y.E. Pokrovsky, Proceedings of the 7th Int. Conf. on Hadron Spectroscopy ‘Hadron’97”, Edited by S.U. Chung and H.J. Wilutzky, AIP, Upton, NY, 1997, p409; Proceedings of the XXXIII Int. Winter Meeting on Nucl. Phys., Edited by I. Iory, Bormio (Italy), 1995, Suppl. **N101**, 540 (1995).
- [12] Th. Walcher, hep-ph/0111279.












This figure "bar191.png" is available in "png" format from:

<http://arXiv.org/ps/nucl-ex/0207004v1>


Calculated (J^P)

Experimental



Spin

Isospin






TABLE I. Experiments whose results are displayed in Fig. 7. The last column (j) of Fig. 7 shows also one mass ($M=1136$ MeV) extracted from the $np \rightarrow pX$ experiment from LAMPF by B.E. Bonner et al., Phys. Rev. D27 497 (1983).

column	reaction	variable	incident energy (MeV)	reference
(a)	$pp \rightarrow \pi^+ pX$	M_X	1520	this work
(b)	$pp \rightarrow \pi^+ pX$	M_X	1805	this work
(c)	$pp \rightarrow \pi^+ pX$	M_X	2100	this work
(d)	$pp \rightarrow \pi^+ pX$	$M_{p\pi}$	1520	this work
(e)	$pp \rightarrow \pi^+ pX$	$M_{p\pi}$	1805	this work
(f)	$pp \rightarrow \pi^+ pX$	$M_{p\pi}$	2100	this work
(g)	$dp \rightarrow ppX$	M_X	1722	this work
(h)	$dp \rightarrow ppX$	M_X	2100	this work
(i)	$\gamma n \rightarrow p\pi^+\pi^0$	$M_{p\pi}$	500-800	(MAMI)
(j)	$\gamma p \rightarrow \pi^+ n$	W_{CM}	300-2000	(Bonn)